



Bernard V. Jackson

A. Buffington

P.P. Hick






CASS/UCSD


HELIOSPHERIC TOMOGRAPHY **Motivation 1: The Helios** Spacecraft Photometer Systems

HELIOS spacecraft: the three photometers are shown as tubes with blackened ends.




HELIOSPHERIC TOMOGRAPHY Helios 2 Spacecraft Contour Image of the May 24 CME

HELIOS 2 spacecraft contour image. May 24, 1979 CME and Video -->



Solwind coronagraph difference image on May 24, 1979.



Motivation: The Solar Mass Ejection Imager - SMEI


SMEI has been delivered to the Coriolis spacecraft for integration. Launch, next year. AF / NASA Funding.



Motivation: NASA's Solar Probe Heliospheric Imager


Proposal to orbit the Sun to within 4 solar radii of its surface.






Heliospheric Imager Design

Solar Orbiter heliospheric imager schematic.



HELIOSPHERIC TOMOGRAPHY Heliospheric Imager Prototype

Solar Orbiter heliospheric imager optical prototype.



Solar Orbiter heliospheric imager optical prototype Images
















Current Heliospheric C.A.T. Analyses









Current Heliospheric C.A.T. Analyses

Line of sight "crossed" components on a reference surface. Corotational projections on the reference surface are shown.





Time-Dependent Tomographic Analysis Technique Reconstruction






image on May 7, 1979.




CASS/UCSD 1979/05/10 12

Time-Dependent Tomographic Analysis Technique Reconstruction




correlation 0.733

April 26,1979 June 6, 1979 Helios
2 video and density
time series
comparison timedependent model
with Helios 2
spacecraft in situ
observations.




Time-Dependent Tomographic Analysis Technique Reconstruction

Helios 1 and 2 Carrington
Rotation 1653 density
reconstruction video. The
Time series correlation is
0.86 for the time-dependent
model comparison with
Helios 2 spacecraft in situ
observations for the
duration of the video.




HELIOSPHERIC TOMOGRAPHY Current IPS Heliospheric Analysis (http://stesun5.stelab.nagoya-u.ac.jp/index-e.html)


STELab IPS array near Fuji (1500 m²)




Time-Dependent Heliospheric C.A.T.


Analysis






Time-Dependent Heliospheric C.A.T. Analysis









Future

- •1000 times more data than Helios!
- •10 times the resolution in all coordinates
- •at least 10⁴ times the computing needs!!





CASS/UCSD 1979/05/10 12

HELIOSPHERIC TOMOGRAPHY Summary

We are currently able to do a pretty accurate job using our time-dependent tomography technique and Helios photometer observations.

Heliospheric imagers SMEI and others will provide nearly 1000 times more data than Helios. This will provide density reconstruction with an ~1-hour time cadence and 1° by 1° latitude - longitude resolution IF the means can be found to analyze the data (real-time?).